


After Medians Comes Nedians

On the day I wrote my last [blog about some interesting properties of the medians](#) of a triangle, I received a package of old Mathematics Teacher articles from Dave Renfro. One of the first I looked at was a January 1951 "Mathematical Miscellanea" edited by Phillip S. Jones. The article contained a contribution by John Satterly of the University of Toronto on a type of cevian that he called "Nedians". [My personal choice, since the "med" root is for the middle, would have just been to call them n-dians, but I'm sure that would have had a cultural backlash.]

For those who may not be familiar with the term "cevian", it refers to a segment in a triangle from a vertex to the opposite side (extended if necessary). Angle bisectors, medians, and altitudes are all cevians then, but a perpendicular bisector of a side would not be because it doesn't necessarily pass through a vertex. The name is in honor of Giovanni Ceva and was originated in France in 1888 and has spread from there.

Professor Satterly seems to have created the term "nedians" as a comparison for medians to describe a cevian that cuts the opposite side $1/n$ th of the way from one vertex to the next. [I shall use the notation 4-nedian for a nedian that cuts $1/4$ th of the way along the opposite side,] A median would be the



2-nedian.

In the image, triangle ABC has 3-nedians AD, BE, and CF where D is $1/3$ of the way from B to C; E is $1/3$ of the way from C to A, etc.

The intersections of the three Nedians of a triangle will form another triangle at their three points of intersection, called the nedian triangle. [JKL in the image].

Professor Satterly seems to have discovered several properties of the nedians, and their nedian triangles which I will give here; and then I have come up with several interesting properties of my own about them that I will add to this blog.

It is not too difficult using affine properties of a triangle to verify many of these.

Professor Satterly showed that the sum of the squares of the nedians would be $\frac{n^2 - n + 1}{n^2}$ times the sum of the squares of the sides of the original triangle. Notice that when $n=2$, this reduces to $3/4$, the ratio given and proved for the medians in [the previous blog](#).


Students may wish to explore these properties by creating Sketchpad or Geogebra interactive models to confirm them, and then challenge themselves to prove them. Proving them is easier with a property of affine geometry. Every triangle is affine equivalent to every other triangle in the plane. Under affine transformations areas and lengths may change, but ratios of them are preserved... which means that you can choose any triangle... a right triangle or an equilateral triangle, to find a property about ratios of lengths or areas, and it will apply to any other triangle... and that is a BIG idea to lock away (I am not well schooled in the particulars of affine geometry, so if I have mis-stated that in some way, please advise).

$$\frac{(n - 2)^2}{n^2 + n - 1}$$

Professor Satterly also stated that the Nedian triangle will have an area of $\frac{(n - 2)^2}{n^2 + n - 1}$ times the area of ABC. Note that for the median, or 2-nedian, the area diminishes to zero since the three medians intersect in a single point.

Professor Satterly suggested the term "backward nedian triangle" for a case in which the $1/n$ ratio went in the opposite order (Let D be $1/3$ of the way from C to B instead). This can be eliminated if we simply allow any real number for the coefficient. Then the backward 3-nedian is just a $3/2$ -nedian in the regular order, and it seems that all his properties are still preserved. Notice that the areas of the 3 and $3/2$ -nedians are equal, but they are not congruent.

Exploring these constructions a little more, I came up with a few more properties that were not in the article. For example, the perpendicular distance from the three vertices of the nedian triangle to any side of ABC will equal the altitude of ABC to that same side. [I call these the sub-altitudes.]



Here the altitude is shown in bold red, and the three corresponding sub-altitudes are shown in dotted red. In the 3-nedian shown the distance from J to side B plus the distance from K to side B plus the distance from L to side B will equal the altitude from B to side B. A similar result exists for each altitude of the triangle. In addition, the three sub-altitudes will always partition the altitude in the same way. The shortest sub-altitude will be

$$\frac{(1)}{n^2 - n + 1} h$$

$$\frac{(n - 1)^2}{n^2 - n + 1} h$$

It is also clear from the last statement that each of the three small triangles at the vertices of A, B, and C will be congruent. Their bases will each be $1/n$ of a base of the original triangle and their heights are

$\frac{(1)}{n^2 - n + 1}h$ so each of them is an equal fraction, $\frac{1}{n^3 - n^2 + n}$ of the original area of ABC. By similar reasoning we see that all three of the quadrilaterals will also have the same area.

I also observed that that each median is partitioned into three parts whose lengths, in order from the

vertex to the opposite side, are $\frac{n}{n^2 - n + 1}$, $\frac{(n - 1)^2 - 1}{n^2 - n + 1}$, and $\frac{1}{n^2 - n + 1}$.

For the 3-medians in the image, for example, the median CF is partitioned so that CJ is 3/7 of CF; JL is 3/7 of CF; and LF is 1/7 of LF. A similar partition holds for the other two 3-medians AD and BE. In a 4-median, the partitions would be 4/14, 8/13, and 1/14.

I'm gonna call that a good days work...please advise of typo's or just plain bad math... and thanks Dave, for another stimulating journal article.